

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of quinine at suitable intervals. The same practice is insisted upon in the public services whose corps are exposed to paludal influences. It is also proposed to make a gratuitous distribution of quinine among indigent persons living in dangerous districts. The idea, aside from the personal benefit to the afflicted individuals, has a wider scope in public hygiene. Any measure calculated to restore to health persons in whose blood malarial organisms exist, or to prevent healthy persons from becoming affected, diminishes the chances of mosquitoes renewing the infection. Thus the spread of malaria is in a degree checked. The public control of quinine, aside from assuring freedom from adulteration and giving public sanction to the use of the drug, is likely to be a considerable source of revenue, because of the fact that in many parts of Italy, quinine is as great a necessity as its proposed commercial associate salt, and quite as much in habitual use as tobacco.

Respectfully,

J. M. EAGER,

Passed Assistant Surgeon, U.S. M. H.S.

The SURGEON-GENERAL,

U. S. Marine-Hospital Service.

JAPAN.

Report from Yokohama.

YOKOHAMA, JAPAN, January 27, 1902.

SIR: I have the honor to transmit herewith an abstract of the bills of health issued at this port for the week ended January 25, 1902. There were 4 vessels inspected with 314 crew, 33 cabin, and 262 steerage passengers. Of the steerage passengers 186 were bathed and their baggage was disinfected. No quarantinable disease has been reported during the week.

Respectfully,

J. Spencer Hough.

Acting Assistant Surgeon, U.S. M. H. S.

The SURGEON-GENERAL,

U. S. Marine-Hospital Service.

MEXICO.

Report from Vera Cruz.

VERA CRUZ, MEXICO, February 9, 1902.

SIR: I have the honor to make the following report for the week

ended February 8, 1902:

From all causes there were reported 40 deaths, 4 of which were from yellow fever. Four cases of yellow fever were reported for the same period. Two deaths were reported from pernicious fever and 6 from tuberculosis.

Respectfully,

S. H. Hodgson,

Acting Assistant Surgeon, U.S. M. H.S.

The SURGEON-GENERAL,

U. S. Marine-Hospital Service.